

HEDIS® Measurement Year 2023 PPO Annual Report

Care Provided to BCBSIL PPO Members		BCBSIL PPO HEDIS Rate	Quality Compass National Average (PPOs and EPOs)
Effectiveness of Care: Prevention and Screening			
• Breast Cancer Screening (BCS-E)		75%	73%
• Cervical Cancer Screening (CCS)		77%	72%
• Colorectal Cancer Screening (COL) - Total		61%	59%
• Chlamydia Screening in Women (CHL) - Total		46%	45%
• Immunizations for Adolescents (IMA) - Combination 2		37%	29%
Effectiveness of Care: Respiratory Conditions			
• Appropriate Testing for Pharyngitis (CWP) - Total		86%	79%
• Pharmacotherapy Management of COPD Exacerbation (PCE)		79%	76%
○ Systemic Corticosteroid		85%	83%
○ Bronchodilator			
• Asthma Medication Ratio (AMR) - Total		88%	86%
Effectiveness of Care: Cardiovascular Conditions			
• Statin Therapy for Patients with Cardiovascular Disease (SPC)			
○ Received Statin Therapy (Total)		86%	82%
○ Statin Adherence 80% (Total)		81%	82%
Effectiveness of Care: Diabetes			
• Kidney Health Evaluation for Patients With Diabetes (KED)			
○ Kidney Health Evaluation for Patients With Diabetes (Total)		45%	43%
• Statin Therapy for Patients with Diabetes (SPD)			
○ Received Statin Therapy		64%	62%
○ Statin Adherence 80%		73%	78%
Effectiveness of Care: Behavioral Health			
• Antidepressant Medication Management (AMM)			
○ Effective Continuation Phase Treatment		63%	64%
• Follow-Up Care for Children Prescribed ADHD Medications (ADD)			
○ Continuation & Maintenance Phase		48%	48%
• Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM)			
○ Blood Glucose and Cholesterol Testing (Total)		37%	36%
• Follow-Up After Hospitalization for Mental Illness - 7 days (Total) (FUH)		58%	46%
• Follow-Up After Emergency Department Visit for Mental Illness - 7 days (Total) (FUM)		56%	47%
• Follow-Up After Emergency Department Visit for Substance Use - 7 days (Total) (FUA)		24%	21%
• Follow-Up After High-Intensity Care for Substance Use Disorder - 7 days (Total) (FUI)		43%	47%

HEDIS® Measurement Year 2023 PPO Annual Report

(Continued from Page 1)

Care Provided to BCBSIL PPO Members		BCBSIL PPO HEDIS Rate	Quality Compass National Average (PPOs and EPOs)
Effectiveness of Care:		Overuse/Appropriateness	
• Appropriate Treatment for Upper Respiratory Infection (URI) - Total		84%	82%
• Avoidance of Antibiotic Treatment for Acute Bronchitis/Bronchiolitis (AAB) - Total		53%	51%
• Appropriate Use of Imaging Studies for Low Back Pain (LBP)		71%	73%
• Avoiding Opioids at High Dosage (HDO)		97%	96%
• Avoiding Opioids from Multiple Prescribers and Multiple Pharmacies (UOP)		99%	99%
• Avoiding potentially risky Continued Opioid Use (COU) >=31 Days (Total)		98%	98%
Access/Availability of Care:			
• Initiation and Engagement of Substance Use Disorder Treatment (IET) <ul style="list-style-type: none"> ○ Engagement of SUD Treatment - Total 		17%	14%
• Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics (APP) - Total		76%	60%

Data based on Measurement Year (MY) 2023.
Results are rounded to the nearest percentage.

HEDIS® is a registered trademark of the National Committee for Quality Assurance (NCQA).